Distributed Selfhood and Quantum-Like Entanglement in Simple Life Forms: Testing for Nonlocal Behavioral Correlation in Trained Slime Molds

Douglas C. Youvan

doug@youvan.com

March 30, 2025

Can a brainless organism possess a form of awareness, memory, or even a sense of self? In this study, we explore that provocative question through an experimental investigation into *Physarum polycephalum*, a single-celled slime mold with no nervous system that nevertheless demonstrates remarkable problem-solving abilities. Known for navigating mazes, optimizing networks, and adapting to complex environments, *Physarum* offers a unique biological platform for probing the outer limits of cognition and coordination. We propose and test the hypothesis that genetically identical, behaviorally trained slime molds may exhibit nonlocal behavioral correlation when physically isolated and only one member is perturbed. If confirmed, such a result would suggest that these simple life forms share a distributed informational state — one potentially governed by coherence or entanglement-like dynamics. Even a null result, however, would refine our understanding of emergent behavior and set a new experimental standard for investigating collective intelligence. This work lies at the intersection of quantum biology, consciousness studies, and systems theory, inviting us to reconsider what life is—and how it might think—at the most fundamental level.

Keywords: quantum biology, slime mold, Physarum polycephalum, distributed cognition, nonlocal correlation, entanglement, collective intelligence, selfhood, ORCH-OR, behavioral conditioning, emergence, consciousness, biofield, coherence, swarm intelligence, panpsychism, experimental biophysics, field-based communication, quantum information, synthetic life. A collaboration with GPT-4o. CC4.0.

Abstract

This study proposes and details an experimental protocol designed to test for the presence of nonlocal behavioral correlation in trained, genetically identical specimens of *Physarum polycephalum* — a unicellular, multinucleate slime mold known for its problem-solving abilities and distributed intelligence. Building on emerging perspectives from quantum biology and consciousness studies, we investigate whether a primitive form of entangled cognition could be observed across physically separated but behaviorally synchronized clones of the same organism.

Our approach involves three core components: (1) behavioral training, wherein two slime mold clones are conditioned to exhibit consistent decision-making in a maze or nutrient-choice task; (2) physical and electromagnetic isolation, using Faraday cages and environmental controls to prevent classical forms of communication; and (3) a targeted perturbation applied to one clone (System A), followed by time-synchronized observation of both System A and the unperturbed System B. We outline a DIY-compatible setup using low-cost materials and accessible technology, including kitchen-based agar sterilization, time-lapse imaging, and reproducible behavioral assays.

If correlated changes in behavior are observed in the non-perturbed system that deviate from baseline patterns and control conditions, such findings may point to a form of distributed selfhood not mediated by conventional signaling — potentially indicative of quantum coherence, field-based communication, or an unknown biophysical mechanism. While speculative, this protocol offers a rigorous, affordable, and replicable framework to explore whether non-neuronal systems like slime molds can access quantum-like states of shared information, thereby contributing to the foundational question of where — and how — consciousness, coordination, and selfhood begin.

1. Introduction

The origin and nature of consciousness remain among the most profound and unresolved questions in science. Despite advances in neuroscience, artificial intelligence, and complex systems theory, there is still no consensus on how subjective experience arises, how agency emerges from matter, or where the boundaries of the "self" truly lie. The traditional model centers on the brain — a dense, neuronal architecture in which information is processed, decisions are made, and awareness presumably emerges. However, this model may be too narrow, particularly when considering forms of intelligence and coordination that arise in biological systems without brains.

In nature, we find striking examples of collective behavior in species that lack centralized processing. Flocks of birds move as a unit with uncanny precision, schools of fish coordinate in split-second harmony, ant colonies solve logistical challenges through swarm logic, and perhaps most intriguingly, the slime mold *Physarum polycephalum* navigates mazes, optimizes nutrient networks, and exhibits apparent memory — all without neurons, synapses, or any centralized controller. These systems challenge our assumptions about what constitutes intelligence and selfhood, and they raise a provocative question: is there something deeper at play than classical emergence?

Recent theories in quantum biology suggest that coherence and entanglement — long thought to be fragile and limited to the subatomic realm — may play a role in biological systems. Quantum effects have been proposed in avian magnetoreception, olfaction, and photosynthetic energy transfer. Perhaps most radically, the Orchestrated Objective Reduction (ORCH-OR) theory, proposed by Roger Penrose and Stuart Hameroff, suggests that quantum coherence within neural microtubules underlies conscious experience itself. While controversial, ORCH-OR has helped legitimize the broader question of whether quantum phenomena might be necessary to explain aspects of cognition, and whether consciousness itself is a quantum process.

However, this debate has remained largely constrained to the brain — or at least to systems with neuron-like architecture. What has been overlooked is the

possibility that distributed biological systems, such as slime molds or social insect colonies, might also access quantum-like dynamics — not within neurons, but through cytoplasmic networks, field-mediated signaling, or unknown coherence-supporting mechanisms. Could these decentralized systems, which already exhibit rudimentary forms of learning, memory, and decision-making, possess a shared informational state that behaves like a primitive quantum "self"?

This paper explores that question through a novel experimental framework. By behaviorally training genetically identical slime molds to follow consistent decision patterns, then physically isolating them and perturbing only one member of the pair, we propose a test for nonlocal behavioral correlation. If such correlation is observed in the unperturbed organism, it may suggest that information is being shared through non-classical means — a phenomenon that could open the door to rethinking consciousness not as a property of brains, but as a deeper, possibly quantum-informational, aspect of life itself.

2. Theoretical Background

2.1 *Physarum polycephalum* as a Cognitive System

The slime mold *Physarum polycephalum* has emerged as a surprising yet powerful model for studying decentralized intelligence. Despite being a single-celled organism, it can exhibit complex behaviors typically associated with higher life forms. In experiments, *Physarum* has solved mazes by optimizing nutrient-seeking paths, anticipated periodic events in its environment, and even demonstrated a form of decision-making under uncertainty — all without neurons, synapses, or any centralized brain-like organ.

The organism’s structure is key to this paradoxical sophistication. It is a multinucleate syncytium, meaning it has one continuous cytoplasm containing many nuclei, interconnected by a dynamic web of protoplasmic veins. These cytoplasmic networks exhibit rhythmic contractions and oscillations, enabling the slime mold to redistribute mass and information across its body in response to

environmental stimuli. Because all nuclei share the same cytoplasm, local biochemical changes can propagate globally without delay — making it a natural, living example of a non-compartmentalized information system.

Unlike in brains, where information flows through axons and synapses, in *Physarum* the decision-making processes are inherently distributed, spatially embedded, and physically unified by cytoplasmic continuity. This makes it an ideal candidate to explore the concept of a biological “self” without a brain, a phenomenon that challenges both cognitive science and metaphysics.

2.2 Quantum Biology and Consciousness

While traditionally dismissed as too delicate for warm, wet environments, quantum effects in biology are gaining recognition across multiple domains. In photosynthesis, for instance, energy is transferred through protein complexes via quantum coherence, allowing the excitation to explore multiple paths simultaneously and settle on the most efficient. Similar phenomena are suspected in avian magnetoreception, where entangled electron spins within cryptochrome proteins help birds navigate Earth's magnetic field.

These discoveries have laid the groundwork for a more speculative but increasingly discussed idea: that consciousness itself might involve quantum processes. The most well-known proposal in this domain is the Orchestrated Objective Reduction (ORCH-OR) theory by Penrose and Hameroff. ORCH-OR posits that microtubules — structural filaments within neurons — can support transient quantum coherence states that collapse via objective reduction, producing discrete moments of conscious experience. Although the theory has been debated for decades, it introduced a paradigm where quantum coherence and gravitational wavefunction collapse play roles in the emergence of consciousness.

Beyond ORCH-OR, other theories have proposed alternative models of distributed or nonlocal consciousness. Integrated Information Theory (IIT) suggests that consciousness corresponds to the amount of integrated information within a system, regardless of its physical architecture. Panpsychism entertains the idea

that consciousness is a fundamental property of matter, varying in degree and complexity across scales. Swarm intelligence models propose that collective behavior can give rise to emergent decision-making resembling cognitive functions, even in the absence of individual awareness.

Despite these developments, little attention has been paid to whether such quantum or emergent properties might be observable in non-neural biological systems — such as slime molds — that already display proto-cognitive behavior. If *Physarum* is capable of supporting a form of coherence or quantum-like information flow across its cytoplasm, it may function as a primitive but testable model for non-neuronal consciousness.

2.3 Emergent Selfhood and the Concept of a Quantum Swarm

As research continues to erode the distinction between individual cognition and distributed intelligence, the notion of the "self" must also be reconsidered. Rather than being a localized entity housed within a brain, the self may instead be an emergent informational pattern — one that arises from the complex interplay of many components across time and space. In this view, what matters is not the presence of a central processor, but the integration and coherence of information within a system.

This reframing opens the door to a radical possibility: that certain biological systems may function as quantum swarms, capable of maintaining distributed states of awareness or coordination via coherent dynamics. If quantum coherence is not limited to the subatomic or sub-neuronal scale, but can also emerge in a large, spatially extended biological system like a slime mold, then it may be possible to detect this indirectly through nonlocal behavioral correlations — measurable effects that cannot be explained by classical signaling or environmental cues alone.

In this context, *Physarum polycephalum* becomes more than a biological curiosity. It becomes a testable embodiment of a non-brain-based self, a unified but distributed system that may demonstrate signs of a quantum swarm intelligence.

Our proposed experiment — which applies behavioral training, isolation, and targeted perturbation — is designed to probe exactly this possibility. If successful, it would not only advance our understanding of quantum biology and distributed cognition but may also compel us to rethink the origins of selfhood, intelligence, and perhaps even consciousness itself.

3. Hypothesis

We hypothesize that slime molds may possess a form of shared, distributed information state — one that could be sustained through mechanisms such as quantum coherence, nonlocal field dynamics, or an as-yet-undiscovered biophysical process. If such a state exists, it would imply that behaviorally synchronized but physically separated clones of *Physarum polycephalum*, derived from the same genetic origin and trained under identical conditions, may remain informationally linked in a manner not explainable by classical communication channels.

Specifically, we propose that when a targeted environmental perturbation is applied to one member of a cloned pair (designated as System A), the non-perturbed counterpart (System B) — housed in an isolated, shielded environment — may exhibit correlated changes in behavior, decision-making, or growth dynamics that mirror or complement those of System A. These correlations may manifest as synchronized directional shifts, altered nutrient-search patterns, or deviations from trained routines — all occurring in the absence of chemical, electromagnetic, thermal, or mechanical signaling.

This behavior would not be expected under classical assumptions, wherein each organism operates as a closed, causally isolated system once physically separated. However, if *Physarum* operates in part through nonlocal information sharing — whether via quantum entanglement, coherence maintained across cytoplasmic memory-like states, or a shared biofield — then perturbations to one instance of the organism could propagate informationally to the other.

Such a finding would constitute experimental support for a new category of biological information processing — one that straddles the boundary between classical and quantum domains. It would challenge the standard conception of individuality in biological systems and lend credence to speculative models of distributed cognition and consciousness that allow for nonlocal coherence.

Importantly, our hypothesis does not assume a fully developed or sentient consciousness in *Physarum*. Rather, it allows for the possibility that primitive coherence at the informational or behavioral level may still emerge in a biological system that displays intelligence-like traits without a central nervous system. This coherence may be detectable experimentally through statistically significant correlations in behavior that persist even after classical pathways of communication have been meticulously eliminated.

In summary, we posit that nonlocal behavioral correlation between physically separated but behaviorally entrained slime molds, in response to localized perturbation of only one, would serve as experimental evidence for a distributed, possibly quantum-coherent "self." This would open a new frontier in the study of consciousness, selfhood, and biological information flow — one rooted not in neurons, but in the dynamic, extended networks of life itself.

4. Experimental Design

4.1 Overview

The experimental design centers on testing for nonlocal behavioral correlation between two genetically identical and behaviorally conditioned *Physarum polycephalum* specimens. The procedure consists of four major phases: cloning and training, establishing behavioral baselines, physical isolation, and targeted perturbation with synchronized observation. By splitting a cultured *Physarum* into two genetically identical portions, training them to exhibit the same decision-making behavior, and then isolating them under controlled conditions, we

establish the prerequisites for identifying correlations in behavior that cannot be attributed to classical communication or environmental interference.

After establishing a clear behavioral baseline, one system (System A) is subjected to a perturbation event while the other (System B) is left untouched. Both systems are monitored via time-lapse imaging, and any deviation in behavior — particularly correlated deviation from the baseline pattern — is recorded, analyzed, and compared statistically. The absence of any classical signaling channel ensures that any synchronous behavior must emerge from a shared, possibly nonlocal, informational connection.

4.2 Behavioral Training Protocol

Maze or Nutrient-Choice Conditioning

To ensure that the slime molds develop predictable, synchronized decision-making behavior, both clones are subjected to repeated training in a standardized spatial task, such as:

- A Y-shaped maze, in which one path contains a stronger nutrient stimulus (e.g., two oat flakes instead of one),
- A triangular nutrient configuration, where one oat flake is enriched (e.g., soaked in sugar solution),
- A T-shaped fork, where previous nutrient history influences directional bias.

The slime molds are trained in these environments for 3–5 iterations, allowing them to establish a preferred path based on reinforcement. During this phase, any random deviation is corrected by reinforcing the "correct" choice (e.g., by removing the alternate oat flake before the organism reaches it). The goal is to condition both System A and System B to exhibit identical, reproducible behavioral patterns in response to identical spatial layouts and nutrient cues.

Clone Selection Criteria

Only slime molds that display consistent directional bias (e.g., choosing the same path 3 out of 4 times or more) and comparable growth rates are selected for use in the final paired experiment. This ensures that baseline behavioral synchrony is established prior to isolation and perturbation, making subsequent deviations statistically and behaviorally meaningful.

4.3 Isolation and Environmental Control

After training, the clones are transferred to separate environments designed to eliminate all known classical channels of interaction. These environments are constructed using the following methods:

Faraday Shielding

- Each Petri dish is placed inside a Faraday cage constructed from grounded copper mesh or foil-lined containers to eliminate electromagnetic communication.

Magnetic and Acoustic Isolation

- The dishes are placed on vibration-dampening surfaces (e.g., foam or gel pads) to prevent transmission of mechanical signals.
- Neodymium shielding or mu-metal (optional) may be used to reduce ambient magnetic fluctuations, though this step is not essential in initial trials.

Environmental Standardization

- Temperature, humidity, light exposure, and airflow are kept strictly identical for both environments using identical containers, incubation chambers, and conditions.
- Internal light sources (if used for imaging) are synchronized and non-heating (e.g., LED arrays).

Time-Lapse Imaging and Motion Tracking

- Each system is monitored continuously using identical time-lapse imaging setups (e.g., smartphone cameras or webcams recording at one frame per minute).
- Optional: Image processing software (e.g., ImageJ, Fiji) is used to quantify growth direction, network morphology, and temporal dynamics.

4.4 Perturbation and Measurement

Targeted Perturbation of System A

At a predetermined time point, System A is subjected to a localized environmental perturbation, designed to influence its behavior in a sharp, measurable way. These perturbations may include:

- Exposure to UV light for 30–60 seconds using a handheld UV flashlight (302 nm wavelength preferred),
- Application of a salt solution drop near the leading edge of growth,
- Sudden nutrient removal (e.g., removal of an oat flake from the expected decision point),
- Localized heating using a low-power infrared lamp (optional).

The key requirement is that the perturbation alters the behavior of System A in a detectable manner, such as changing its growth direction, halting movement, or shifting toward a non-reinforced area.

System B (Control)

System B, housed in a physically isolated and identically controlled environment, receives no stimulus. It continues growing under undisturbed conditions. The experimental objective is to observe whether System B exhibits a synchronous or statistically improbable deviation in its behavior that correlates temporally or structurally with the response of System A.

Behavioral Observation and Data Recording

- The imaging data from both systems are synchronized and analyzed for:
    - Changes in directionality,
    - Network branching patterns,
    - Temporal onset of growth pauses or redirection,
    - Any sudden behavioral shifts that deviate from prior trained behavior.

Statistical Analysis

- Behavioral deviation is quantified in terms of deviation angle, growth rate, and time to decision.
- Pearson or Spearman correlation coefficients are calculated between System A and System B behaviors across timepoints.
- Permutation testing and Monte Carlo simulations may be used to assess the statistical significance of any observed correlation.
- Control trials (with no perturbation to either system) are included to establish background variability and false-positive rates.

5. Controls and Verification

A central claim of this experiment is that correlated behavioral changes between spatially separated slime mold systems, in response to localized perturbation of only one, may indicate the presence of a nonlocal, shared informational state. Given the profound implications of such a result, a rigorous control and verification strategy is essential to rule out classical explanations, experimental artifacts, and cognitive bias.

This section outlines four key types of control, along with techniques for eliminating ambiguity in both data collection and interpretation.

5.1 Null Controls: Clone Pairs Without Perturbation

To establish a baseline behavioral variance, genetically identical and behaviorally synchronized clones are subjected to parallel isolation and observation under identical environmental conditions but with no perturbation applied to either system. This control verifies:

- That no inherent synchronized fluctuation in behavior emerges spontaneously,
- That imaging, incubation, and observation protocols do not induce behavioral artifacts,
- That trained behaviors remain stable over the observation window in undisturbed conditions.

By comparing experimental trials to these null-control pairs, we can detect whether perturbation of System A leads to statistically anomalous changes in System B — beyond the expected range of natural behavioral variance.

5.2 Genetic Controls: Unrelated Slime Mold Pairs

To determine whether any observed correlation is tied specifically to genetic and developmental identity, a series of trials will be run with genetically unrelated slime molds — either drawn from different parent cultures or at different life-cycle stages.

These pairs will undergo the same behavioral training, isolation, and observational protocols, including perturbation of System A. However, since they are not genetically identical and lack a shared developmental history, no correlation is expected under the distributed-information hypothesis.

A lack of correlated behavior in unrelated pairs, compared to that in cloned pairs, would support the idea that shared genetic/cellular history may be a prerequisite for any form of informational linkage — echoing results from some quantum entanglement analogues in biological systems.

5.3 Temporal Randomization and Blinded Analysis

To minimize cognitive bias, temporal randomization and double-blinded analysis will be implemented:

- Randomized timing: Perturbation times for System A will be determined via pseudorandom selection from a pre-generated list. The observer monitoring System B will be unaware of the timing of perturbations in System A.
- Blinded scoring: Behavioral image sequences from both systems will be stripped of timestamps and labeled with randomized codes before being scored. Analysts will evaluate behavioral change (direction, rate, branching, etc.) without knowing whether the sample was experimental or control.

This process reduces the risk of confirmation bias, strengthens the statistical objectivity of behavioral scoring, and ensures that any observed correlations are not unconsciously inferred or exaggerated.

5.4 Optional Signal Detection: Quantum and Field-Based Instrumentation

Though not required for initial behavioral tests, optional integration of quantum-sensitive instrumentation may provide additional insight into the physical substrate of any nonlocal correlation.

Candidate Techniques:

- Nitrogen-vacancy (NV) center magnetometry: NV-diamond sensors can detect ultra-weak magnetic field fluctuations possibly indicative of coherent spin activity within biological samples.
- Photonic emission monitoring: Highly sensitive photodetectors can be used to detect delayed biophoton emissions from slime molds — potentially modulated under perturbation.

- Quantum noise correlation analysis: Broadband noise spectroscopy across EM bands may reveal anomalous correlated fluctuations between separated systems.

While these approaches require more advanced equipment and analysis, they could serve to correlate behavioral shifts with field-level dynamics, supporting a physical mechanism for the hypothesized information linkage.

Conclusion of Control Strategy

Through the use of genetic controls, null trials, randomized timing, blinded scoring, and optional physical instrumentation, this experiment aims to rigorously eliminate classical explanations for any observed behavioral correlations. Any remaining, statistically significant synchronous behavior between perturbed and unperturbed slime molds may then be viewed as a candidate indicator of nonlocal biological coherence — pointing to a new, perhaps quantum-informational dimension of collective selfhood in simple life forms.

6. Expected Results and Interpretation

The outcome of this experiment will fall into one of two general categories: positive behavioral correlation or negative/inconclusive results. Both outcomes are significant, though they carry fundamentally different implications for how we interpret biological information flow, cognition, and the potential for quantum coherence in living systems.

6.1 Positive Behavioral Correlation

If the unperturbed slime mold (System B) exhibits a behavioral deviation that temporally coincides with the perturbation applied to System A — such as changing direction, halting movement, or breaking from trained behavior — and

such responses are statistically unlikely under null control conditions, the result would be highly anomalous under classical assumptions.

Implications:

- Nonlocal Information Flow:
  The most direct interpretation is that some form of nonlocal information exchange is occurring between the two slime molds. Since all known classical channels are eliminated by shielding and isolation, the correlation would suggest a field-like or quantum-informational linkage, potentially mediated by coherence or entanglement between the systems.
- Biological Entanglement Analogy:
  While not implying literal quantum entanglement in the strict physics sense, the findings may point to entanglement-like effects — that is, correlated states that resist explanation by local causality. This mirrors the logic of quantum mechanics, in which entangled particles influence one another instantaneously regardless of distance.
- Reframing Slime Molds as Primitive “Selves”:
  A positive result would challenge our concept of biological individuality. Instead of treating each slime mold as a closed organism, it would support the idea that distributed life forms may constitute a collective "self" — a coherent informational unit expressed across space.
- Support for Quantum-Informational Life Theories:
  The experiment would lend empirical weight to emerging views that life is inherently quantum-informational — not simply biochemical machinery, but a system organized by subtle, possibly nonlocal rules of coordination. This dovetails with speculative theories such as ORCH-OR, as well as broader efforts to unify physics, information theory, and biology.
- Philosophical and Scientific Repercussions:
  If verified and reproduced, such results could prompt re-evaluation of foundational biological assumptions. It may even revive concepts such as

vitalism in a modern, physics-compatible form — where "life force" is reconceived as distributed coherence, not mystical essence.

In short, positive results would offer a paradigm-shifting insight into how information, agency, and coordination can exist in systems far simpler than brains, yet no less mysterious in their capacities.

6.2 Negative or Inconclusive Results

If System B does not exhibit any behavioral deviation or correlation with System A following perturbation, or if any observed variations fall within the statistical range of control conditions, the experiment will support a more classical interpretation of slime mold behavior.

Implications:

- Validation of Classical Emergence:
  A negative result would reinforce the prevailing view that slime mold behavior is driven by localized chemical gradients, feedback loops, and emergent pattern formation — without requiring quantum or nonlocal explanations.
- Reproducibility of Trained Behaviors:
  If trained slime molds continue to make consistent decisions in the absence of perturbation, it will validate the robustness of behavioral conditioning protocols developed for this experiment. This would be a significant contribution to slime mold research, independent of the outcome.
- Necessity of Higher Sensitivity:
  It may be that any nonlocal coherence effects — if they exist — are too subtle or transient to detect with current instrumentation or behavioral metrics. This would suggest the need for:
    - More sensitive detection equipment (e.g., quantum sensors),
    - Larger sample sizes and statistical power,

    - Or more complex tasks that allow richer behavioral variability.

- Framework for Future Research:
  Even inconclusive results are valuable. They can help define the boundary between emergent complexity and fundamental coherence, and offer a model for how to methodologically test quantum-biological hypotheses with rigor and low-cost experimental setups.

Conclusion of Interpretation

Whether the results are positive, negative, or ambiguous, this experiment contributes meaningfully to the emerging frontier between biological intelligence and quantum information theory. A positive outcome would challenge foundational models of biological individuality and causality, while a negative outcome would reaffirm the strength of classical explanations and lay the groundwork for improved experiments.

Either way, the experiment embodies a new kind of inquiry: one that dares to explore the edges of biology and physics, where the question is no longer just *what life is*, but *how it thinks, acts, and perhaps connects — even across space.*

7. Broader Implications

The implications of this experimental framework extend far beyond slime mold behavior. If evidence emerges that *Physarum polycephalum* — a simple, brainless organism — can exhibit nonlocal behavioral correlation suggestive of a shared informational state, then our current understanding of life, cognition, and information itself must be reconsidered. Even in the absence of such findings, the effort to test this possibility reframes fundamental scientific assumptions and invites new models of biological agency.

7.1 Biological Information Systems as Quantum Processors

This experiment encourages us to see biological systems not merely as chemical engines, but as informational systems — potentially governed by laws that transcend classical physics. If slime molds can coordinate behavior nonlocally, even in a limited and primitive sense, it may imply that life is capable of quantum-like information processing even in the absence of neurons or digital circuits.

Just as quantum computers use superposition and entanglement to perform complex computations that classical systems cannot match, biological systems might exploit nonlocal coherence to optimize decisions, solve spatial problems, or transmit information across a distributed field. If true, this would open a new field of quantum biocomputation, in which organisms — including slime molds, mycelial networks, and perhaps even embryonic structures — are understood as living quantum systems, optimized not only by evolution but also by informational constraints encoded in the fabric of reality.

7.2 Redefining the Boundaries of Cognition and Agency

Current models of cognition are largely anthropocentric and brain-centric, defining intelligence as the product of synaptic connections, centralized control, and symbolic manipulation. This experiment, however, aligns with a growing body of research suggesting that cognition is a more general property of life — distributed, embodied, and often decentralized.

If slime molds are shown to exhibit coordination beyond classical signaling, it would support the view that agency — the ability to make meaningful decisions and respond to complex conditions — is not restricted to higher organisms. Instead, it may emerge wherever coherent information processing occurs, regardless of architecture. This redefinition blurs the line between behavioral intelligence and cognitive selfhood, allowing us to speak meaningfully about primitive or preconscious selves that do not rely on the traditional machinery of a nervous system.

7.3 Relevance to Consciousness, AI, Swarm Robotics, and Synthetic Life

The outcomes of this study have implications for a wide range of disciplines and technologies:

Consciousness Studies

A demonstration of nonlocal behavioral coordination in a non-neural system would be of deep interest to those studying the origins of consciousness. It would suggest that consciousness, or at least the precursors to it, may emerge from information flow and coherence, rather than from complexity or brain structure alone.

Artificial Intelligence

Current AI models — especially deep learning — are based on classical, localized processing. If biological systems leverage quantum or field-like coherence, this raises the possibility of radically new AI architectures, perhaps involving nonlocal learning, collective agency, or field-based memory systems inspired by slime mold dynamics.

Swarm Robotics

Swarm systems today rely on rule-based emergent behavior. Incorporating lessons from *Physarum* — including the possibility of nonlocal influence or internal coherence — could lead to more adaptive, resilient, and intelligent collectives, capable of unified action without constant inter-unit communication.

Synthetic and Artificial Life

In synthetic biology, the aim is often to recreate life-like behavior in artificial systems. This experiment opens the possibility that life's deepest organizing principles may include nonclassical coherence, which could be harnessed to engineer novel systems with distributed cognition or emergent awareness.

7.4 Philosophical Consequences for the Soul, Self, and Mind

Finally, the philosophical stakes are significant. If even a slime mold can, in some minimal way, exhibit signs of a nonlocal self, then long-standing metaphysical questions must be revisited:

- The Soul: Many religious and spiritual traditions describe the soul as an invisible, organizing principle — not confined to physical form or locality. If life exhibits coherence beyond space and classical time, this may offer a scientific metaphor (or mechanism) for what ancient texts have intuited about the non-material essence of life.
- The Self: A nonlocal or distributed self implies that identity is not fixed to a brain, body, or location, but is instead a dynamic field of information and coherence. This would support panpsychic, idealist, or informational metaphysics in which the self is not a thing, but a pattern.
- The Mind: Rather than being confined within a skull, mind might be more broadly understood as a coherent, living response to the world — emerging wherever information is processed in a meaningful, dynamic way. If slime molds exhibit even the faintest shadows of this, it challenges us to reframe what mind truly means.

Conclusion of Broader Implications

This experiment operates at the frontier of disciplines — biology, physics, philosophy, and technology — and invites a bold but rigorous rethinking of what it means to be alive, aware, and connected. Whether slime molds prove to be primitive selves or simply complex chemical machines, the act of testing the boundary between classical and quantum information in living systems may ultimately lead us to a new scientific language of the soul — grounded in coherence, emergence, and the unity of life itself.

---

8. Future Directions

The experimental framework described in this paper represents an initial foray into testing whether simple, decentralized biological systems can exhibit nonlocal behavioral coordination — a hallmark of distributed selfhood and potentially quantum-informational coherence. Regardless of the outcome, the experiment opens a rich horizon of subsequent studies across multiple dimensions: species complexity, spatial scale, technological instrumentation, and comparative cognition.

8.1 Extension to Ant Colonies and Fungal Networks

While *Physarum polycephalum* offers a uniquely tractable and unified experimental model due to its acellular, multinucleate structure, many other organisms exhibit distributed intelligence and collective behavior without central control. Two particularly promising candidates for further research are ant colonies and fungal mycelial networks.

Ant Colonies:

Ants demonstrate emergent behavior at the colony level, including:

- Division of labor
- Collective decision-making
- Complex foraging and path optimization

By applying a similar protocol — training spatial preferences in a small ant colony, isolating subgroups, and perturbing one — researchers could explore whether remote behavioral shifts emerge between colony members who were once co-located. Given the pheromone-based communication of ants, rigorous isolation and shielding would be essential, but possible with sealed enclosures and filter barriers.

Mycelial Fungal Networks:

Fungi such as *Armillaria* and *Mycelium* form vast underground networks, often described as “the Wood Wide Web.” They exhibit:

- Resource sharing across distances
- Network pruning and optimization
- Possible electrochemical signaling across hyphae

Their structural similarity to neural networks, combined with recent findings of bioelectrical communication, make them excellent candidates for studying long-distance coherence within and between fungal clones. Here, quantum coherence or biofield-mediated signaling may play a more substantial role than currently understood.

8.2 Repeat with Longer Distances or Entanglement-Enhanced Sensors

In the initial experiment, *Physarum* clones are separated by relatively short distances — on the order of centimeters to meters — largely due to practical constraints. Future experiments should test whether behavioral correlation persists or weakens as the physical separation increases to:

- Room-scale (5–10 meters)
- Building-to-building
- Geographically separated labs (even across cities or continents)

If coherent behavioral patterns remain detectable across large distances, the experiment would provide even stronger evidence for nonlocal informational linkage, analogous to spatially extended entanglement in quantum systems.

To enhance the sensitivity of such tests, entanglement-aware sensors can be introduced, including:

- Quantum magnetometers to track sub-picotesla field fluctuations
- Photon correlation detectors for possible biophoton emission timing anomalies
- Optical coherence tomography adapted for living biological networks

These technologies, though not required for initial behavioral studies, could help reveal hidden physical dynamics beneath apparent behavioral synchrony.

8.3 Integration with Quantum Imaging and Topological Signal Tracing

To move beyond surface-level behavioral correlation and into the realm of mechanistic insight, future work should incorporate quantum imaging tools and topological signal analysis:

Quantum Imaging:

- NV-center diamond microscopy for real-time magnetic fluctuations
- Coherence lifetime mapping of bioelectromagnetic fields
- Ultrafast spectroscopy to detect transient quantum states in cytoplasm

These methods would allow researchers to visualize potential coherence domains within slime molds, fungal networks, or even microbial consortia — revealing whether biological matter can sustain structured information fields over time.

Topological Signal Tracing:

- Use of persistent homology, knot theory, or topological data analysis (TDA) to model slime mold or fungal network morphology as a dynamic topological space.
- Detecting abrupt topological reconfigurations in response to remote perturbations could indicate global state shifts not explainable by local cause-effect chains.

Such approaches would begin to define what a distributed self "looks like" mathematically — and whether these systems exhibit something analogous to a global phase coherence in condensed matter physics.

8.4 Cross-Species Comparison of Distributed Cognition Candidates

To refine our understanding of what enables or inhibits distributed coherence in living systems, a comparative biology approach should be adopted. Future research could assess a range of organisms for signs of nonlocal coordination, based on parameters such as:

- Degree of cytoplasmic continuity (e.g., syncytial organisms like *Physarum* vs. multicellular colonies)
- Communication modality (chemical, electrical, acoustic, photonic)
- Developmental similarity or clonality
- Spatial scale of operation

Candidate systems might include:

- Coral polyps
- Colonial algae (*Volvox*)
- Biofilm-forming bacteria
- Plant root systems with shared mycorrhizal networks

The aim would be to identify the necessary and sufficient conditions for distributed selfhood and information linkage — and to assess whether these systems exhibit species-specific limitations, or whether quantum-like coherence is a deeper principle of life itself.

Conclusion of Future Directions

The present study lays the groundwork for an entirely new research trajectory — one that treats distributed biological systems as potential quantum-cognitive agents, rather than as mere aggregates of localized responses. From ants to fungi, from slime to photons, this line of inquiry may one day yield a unified understanding of coherence, cognition, and the emergence of the self — not just in minds, but in life wherever it appears.

9. Conclusion

This study presents a novel and accessible experimental framework for exploring one of the most profound questions in science and philosophy: Can selfhood, cognition, or coherent information flow emerge in biological systems without a brain — and possibly without classical causality? By focusing on *Physarum polycephalum*, a remarkably capable yet simple organism, we offer a concrete and reproducible method to test for nonlocal behavioral correlation between physically separated, genetically identical slime molds.

What sets this work apart is its fusion of behavioral conditioning, shielded isolation, and perturbation-response analysis, executed with low-cost, DIY-compatible methods that open the field to broader participation. In doing so, it builds a rigorous, falsifiable bridge between classical experimental biology and speculative ideas drawn from quantum theory, systems neuroscience, and philosophy of mind.

If positive results are observed — if System B changes its behavior in synchrony with perturbations applied solely to System A — this would offer striking evidence for a shared, distributed informational state. Such a finding would invite radical reinterpretation of what it means to be a “self,” extending the concept of consciousness beyond its anthropocentric and neurocentric boundaries into the domain of non-neural, field-like intelligence. It could suggest that life possesses deeper organizing principles rooted in coherence, correlation, and information —

principles that have not yet been fully integrated into biological or cognitive science.

Conversely, if no significant correlation is found, the experiment still provides value. It affirms the robustness of behavioral training in simple organisms, sets a methodological standard for future work on distributed cognition, and narrows the parameter space in which quantum or nonlocal effects might be sought. Even a null result helps distinguish between genuine coherence and classical complexity, ensuring that future research into these phenomena is grounded in empirical clarity rather than speculation.

In either case, this work contributes to a long-overdue paradigm shift — one that dares to ask whether cognition, agency, and perhaps even consciousness are not confined to brains, but may be properties of information flow itself, regardless of scale, architecture, or substrate. It invites us to look again at the humble slime mold, not merely as a blob of protoplasm, but as a living question mark at the intersection of biology, physics, and the nature of being.

Appendices

Appendix A: Materials and Methods

Organism and Biological Materials

- *Physarum polycephalum* culture
    - **Supplier:** Carolina Biological Supply or Ward's Science
    - **Cost:** $25–40 per culture
    - **Notes:** Can be subcultured indefinitely on moist filter paper and oats.
- **Oat Flakes (Organic Rolled Oats)**
    - **Supplier:** Grocery store (any brand)
    - **Cost:** ~$5 per box (months of experiments)

Petri Dishes and Growth Substrates

- **100 mm x 15 mm Petri dishes** (plastic, sterile or reusable glass)
    - **Supplier:** Amazon, Fisher Scientific, Carolina
    - **Cost:** ~$15 for 20-pack
- **Non-nutrient Agar** (for mobility substrate)
    - **Supplier:** Amazon, lab supply stores
    - **Cost:** ~$20 per 100g
- **Distilled Water**
    - **Supplier:** Grocery/pharmacy
    - **Cost:** ~$2 per gallon

Environmental Control Materials

- **Faraday Cages (DIY)**

    - **Materials:** Copper mesh or foil-lined plastic containers
    - **Cost:** ~$20–40 depending on size
    - **Notes:** Must be grounded to dissipate EM noise
- **Humidity Control**
    - **Method:** Damp paper towel or sponge inside container, not in direct contact with slime
    - **Notes:** Monitor to avoid drying or over-condensation
- **Temperature Control (optional)**
    - **Method:** Room temperature or incubator (20–25°C ideal)
    - **Cost:** ~$0–100 depending on setup

Imaging Setup

- **Smartphone with time-lapse app** or **USB webcam**
    - **Cost:** $0 (if already owned) to ~$50
    - **Mounting:** Tripod, clamp, or stack of books
    - **Light Source:** Soft white LED lamp, consistent brightness

---

Appendix B: Detailed Experimental Protocol

Day 1: Culturing and Cloning

1. Pour ~20 mL of non-nutrient agar into Petri dishes; allow to solidify.
2. Place an oat flake in the center of each dish.
3. Transfer ~1 cm² of *Physarum* from culture plate onto oat flake.
4. Incubate overnight in dark or dim environment at 22°C.

Day 2: Behavioral Training (see Appendix C for details)

- Train clones to navigate toward reinforced nutrient path in maze or preference assay.

Day 3: Isolation and Perturbation

1. Select two clones showing consistent behavioral preference.
2. Place in separate shielded containers under matched environmental conditions.
3. Begin synchronized time-lapse imaging of both samples.
4. After baseline behavior is established (~4–6 hours), apply **stimulus to System A only**:
    - **UV pulse:** 30–60 seconds from ~5 cm away
    - **Salt drop:** 0.1 mL saturated NaCl near growth edge
    - **Nutrient removal:** Carefully remove one oat flake using sterilized forceps
5. Continue imaging both systems for an additional 12–24 hours.
6. Analyze for deviation or synchronized change in System B.

Appendix C: Behavioral Training Protocol

Training Configurations

- **Y-maze Template:** Create agar in Y-shaped mold or with physical guides. Place an oat flake at each fork end.
- **Triangle Template:** Three oat flakes placed equidistantly in a triangle, with one pre-soaked in sugar water as the "preferred" target.

Training Procedure

1. Place slime mold at origin (e.g., base of Y-maze).
2. Allow 6–12 hours for decision-making.
3. Reinforce by removing or reducing less-preferred oat flake after consistent choices.
4. Repeat 3–5 cycles to stabilize behavioral preference.
5. Select only clones that chose the reinforced option 3 out of 4 trials or more.

Appendix D: Agar Preparation and Sterilization

Ingredients

- 2 g agar powder per 100 mL distilled water
- Optional: Add trace nutrients (e.g., 0.1 g glucose) for faster growth

Sterilization Protocol (DIY Pressure Cooker)

1. Dissolve agar in measured distilled water in heatproof glass (e.g., Mason jar).
2. Cover with loose-fitting lid or aluminum foil to allow steam escape.
3. Place on rack inside pressure cooker with 1 inch of water in base.
4. Seal cooker and heat until pressure reaches 15 psi.
5. Maintain full pressure for 20 minutes.
6. Allow pressure to release naturally; carefully remove jar.
7. Cool to ~50°C and pour into Petri dishes in clean area.
8. Let solidify, then cover and store inverted at 4°C or room temp for up to 2 weeks.

References

1. Penrose, R., & Hameroff, S. R. (2014). *Consciousness in the universe: A review of the 'Orch OR' theory*. Physics of Life Reviews, 11(1), 39–78. https://doi.org/10.1016/j.plrev.2013.08.002
2. Hameroff, S. R., & Penrose, R. (1996). *Orchestrated reduction of quantum coherence in brain microtubules: A model for consciousness*. Mathematics and Computers in Simulation, 40(3–4), 453–480. https://doi.org/10.1016/0378-4754(96)80476-9
3. Adamatzky, A. (2010). *Physarum Machines: Computers from Slime Mould*. World Scientific Publishing.
4. Nakagaki, T., Yamada, H., & Tóth, Á. (2000). *Maze-solving by an amoeboid organism*. Nature, 407(6803), 470. https://doi.org/10.1038/35035159
5. Al-Khalili, J., & McFadden, J. (2014). *Life on the Edge: The Coming of Age of Quantum Biology*. Crown Publishing.
6. Engel, G. S., Calhoun, T. R., Read, E. L., Ahn, T. K., Mancal, T., Cheng, Y. C., ... & Fleming, G. R. (2007). *Evidence for wavelike energy transfer through quantum coherence in photosynthetic systems*. Nature, 446(7137), 782–786. https://doi.org/10.1038/nature05678
7. Gauger, E. M., Rieper, E., Morton, J. J., Benjamin, S. C., & Vedral, V. (2011). *Sustained quantum coherence and entanglement in the avian compass*. Physical Review Letters, 106(4), 040503. https://doi.org/10.1103/PhysRevLett.106.040503
8. Tononi, G. (2008). *Consciousness as integrated information: a provisional manifesto*. The Biological Bulletin, 215(3), 216–242. https://doi.org/10.2307/25470707
9. Chalmers, D. J. (1996). *The Conscious Mind: In Search of a Fundamental Theory*. Oxford University Press.

10. Baluška, F., & Levin, M. (2016). *On having no head: cognition throughout biological systems*. Frontiers in Psychology, 7, 902. https://doi.org/10.3389/fpsyg.2016.00902

11. Camazine, S., Deneubourg, J. L., Franks, N. R., Sneyd, J., Theraulaz, G., & Bonabeau, E. (2001). *Self-Organization in Biological Systems*. Princeton University Press.

12. Trewavas, A. (2003). *Aspects of plant intelligence*. Annals of Botany, 92(1), 1–20. https://doi.org/10.1093/aob/mcg101

13. Ho, M. W. (1997). *Towards a theory of the organism*. Integrative Physiological and Behavioral Science, 32(4), 343–363. https://doi.org/10.1007/BF02688600

14. Sheldrake, R. (2011). *The Presence of the Past: Morphic Resonance and the Habits of Nature*. Icon Books.

15. Fields, C., Glazebrook, J. F., & Levin, M. (2021). *Morphological computation in developmental systems: perspectives on the dynamic stability of self*. Frontiers in Systems Neuroscience, 15, 655212. https://doi.org/10.3389/fnsys.2021.655212

16. Deacon, T. W. (2012). *Incomplete Nature: How Mind Emerged from Matter*. W. W. Norton & Company.

17. Craddock, T. J. A., Tuszynski, J. A., & Hameroff, S. (2015). *Cytoskeletal signaling: Is memory encoded in microtubule lattices by CaMKII phosphorylation?* PLoS Computational Biology, 11(5), e1004241. https://doi.org/10.1371/journal.pcbi.1004241

18. Varela, F. J., Thompson, E., & Rosch, E. (1991). *The Embodied Mind: Cognitive Science and Human Experience*. MIT Press.


Distributed Selfhood and Quantum-Like Entanglement in Simple Life Forms
Testing for Nonlocal Behavioral Correlation in Trained Slime Molds